

**TECHNIQUES IN NON-ABELIAN ADDITIVE COMBINATORICS,
EXAMPLES SHEET 1**

Lent Term 2016

W. T. G.

General remark: I have not redone all these questions, and therefore some of the bounds I state may be a little approximate. If you get different, but similar bounds, then don't worry!

1. The subset $\{0, 1\}^n$ of \mathbb{F}_3^n has density $(2/3)^n$ and contains no affine line. Can you improve on this bound?

2. (i) For each triple of positive integers m, r and d , let $S_{m,r,d}$ be the set of all points $x = (x_1, \dots, x_d) \in \{0, 1, \dots, m-1\}^d$ such that $\sum_i x_i^2 = r$. For each d and m prove that there exists r such that $|S_{m,r,d}| \geq m^{d-2}/d$.

(ii) Prove that no $S_{m,r,d}$ contains a non-trivial triple (x, y, z) with $x + z = 2y$.

(iii) By considering a suitable projection of a suitable set $S_{m,r,d}$, deduce that for each n there exists a set $A \subset \{1, \dots, n\}$ of density at least $\exp(-c\sqrt{\log n})$ (where c is some absolute constant) that contains no arithmetic progression of length 3.

[The above example was discovered by Behrend in 1947. The bound has stayed almost still since then.]

3. Let A and B be subsets of Abelian groups. We say that they are *Freiman isomorphic of order k* if there is a bijection $\phi : A \rightarrow B$ such that the equation

$$\phi(a_1) + \dots + \phi(a_k) = \phi(a'_1) + \dots + \phi(a'_k)$$

holds for elements of A if and only if the equation

$$a_1 + \dots + a_k = a'_1 + \dots + a'_k$$

holds. Prove that if A and B are subsets of \mathbb{F}_p^n that are Freiman isomorphic of order 8, then $2A - 2A$ contains a subspace of dimension d if and only if $2B - 2B$ contains a subspace of dimension d .

4. Let A be a subset of \mathbb{F}_p^n and suppose that $|8A - 8A| \leq C|A|$. Let X be a subspace of \mathbb{F}_p^n of codimension d , chosen uniformly at random from all such subspaces, and let P_X

be the quotient map from \mathbb{F}_p^n to \mathbb{F}_p^n/X . Prove that as long as $p^d > C$, there is a non-zero probability that $P_X(A)$ and A are Freiman isomorphic of order 8. Use this result to prove that $2A - 2A$ contains a large subspace (with a meaning of “large” that you should work out for yourself).

5. Use the inequality $[A_1, A_2, A_3, A_4] \leq \|A_1\|_{\square} \|A_2\|_{\square} \|A_3\|_{\square} \|A_4\|_{\square}$ to prove the inequality $|\mathbb{E}_{x,y} A(x,y)u(x)v(y)| \leq \|A\|_{\square} \|u\|_2 \|v\|_2$.

6. Let $A : \mathbb{C}^n \rightarrow \mathbb{C}^m$ and let $u \in \mathbb{C}^n$ be such that $\|Au\|_2 = \|A\|_{\text{op}} \|u\|_2$. Prove that if $\langle u, v \rangle = 0$, then $\langle Au, Av \rangle = 0$. Use this result to prove that A has a singular-value decomposition.

7. Use the Cauchy-Schwarz inequality several times to prove an inequality of the form

$$|\mathbb{E}_{x,y,z} f(x,y,z)a(x)b(y)c(z)| \leq \Phi(f) \|a\|_2 \|b\|_2 \|c\|_2,$$

where Φ is a functional that you should determine. (If you get the right one, it is in fact a norm, but you do not need to prove this – though it’s not a bad exercise to do so.)

8. Obtain also an inequality of the form

$$|\mathbb{E}_{x,y,z} f(x,y,z)u(x)v(y)w(z)| \leq \Psi(f) \|u\|_{\infty} \|v\|_{\infty} \|w\|_{\infty},$$

where Ψ is an obvious three-dimensional generalization of the box norm.

9. Let A be a subset of \mathbb{F}_p^n of density α . What can you say about the dimension of the largest affine subspace that is contained in $A + A + A$?

10. Prove that the Bohr set $B(K, \epsilon) \subset \mathbb{Z}_N$ has density at least $(\epsilon/2)^{|K|}$. Prove also that it contains an arithmetic progression of length at least $(\epsilon/2)N^{1/|K|}$.

11. Let $A_r \subset \mathbb{F}_2^n$ be the set of all x such that $x_i = 1$ for at most r values of i . Prove that for suitably chosen r (depending on n) the set A_r has density $\frac{1}{2} - o(1)$ but the set $A_r - A_r$ does not contain a subspace of bounded codimension.